Neuroscience Letters, 209 (1996) 218–220
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Abe, S., see Hori, T., 153
Aitken, A., see Layfield, R., 57
Albanese, V., see Strosznajder, J., 145
Alberghina, M., see Strosznajder, J., 145
Allard, P., see Norlén, M., 161
An, L., see Pan, J.C., 121
Andersson, J., see Imamura, K., 185
Apfelbaum, J.L., see Zacny, J.P., 73

Baba, A., see Hori, T., 153
Battaglioli, G., Martin, D.L., Glutamine stimulates γ-aminobutyric acid synthesis in synaptosomes but other putative astrocyte-to-neuron shuttle substrates do not, 129
Beck, O., see Stiller, C.-O., 165
Benoit, G., see Giuliano, F., 1
Bergquist, J., see Stiller, C.-O., 165
Bhatti, T., see Seifritz, E., 41
Blödorn, B., Mäder, M., Urade, Y., Hayaishi, O., Felgenhauer, K., Brück, W., Choroid plexus: the major site of mRNA expression for the β-trace protein (prostaglandin D synthase) in human brain, 117
Blomqvist, A., see Hallbeck, M., 125
Boda, B., Szente, M., Nitric oxide synthase inhibitor facilitates focal seizures induced by aminopyridine in rat, 37
Borst, J.G.G., Lodder, J.C., Roubos, E.W., Kits, K.S., In situ recordings of presumed folliculo-stellate cells in the intermediate lobe of the pituitary gland of *Xenopus laevis*, 61
Brodin, E., see Stiller, C.-O., 165
Brown, K., see Giuliano, F., 1
Brück, W., see Blödorn, B., 117
Bulleit, R.F., Lin, X., Jeppi, J.V., A novel gene selectively expressed in the cerebellum, 93
Burazin, T.C.D., Gundlach, A.L., Rapid but transient increases in cholecystokinin mRNA levels in cerebral cortex following amygdaloid-kindled seizures in the rat, 65

Carfagna, N., Cavanus, S., Damiani, D., Salmoiraghi, P., Fariello, R., Post, C., Modulation of phosphoinositide turnover by chronic nicergoline in rat brain, 189
Cash, C.D., Hechler, V., Mersel, M., Maitre, M., Kinetic characterisation and solubilisation of γ-hydroxybutyrate receptors from rat brain, 25
Cavanus, S., see Carfagna, N., 189
Celio, M.R., see Wintergerst, E.S., 173
Cervero, F., see Herrero, J.F., 21
Chalimoniuk, M., see Strosznajder, J., 145
Chang, S.L., see Quercia, S., 89
Chen, A.C.N., Bragard, D., Plaghki, L., Direct isolation of ultra-late (C-fibre) evoked brain potentials by CO_2 laser stimulation of tiny cutaneous surface areas in man, 81
Chen, A.C.N., see Bragard, D., 81
Cho, A.M., see Zacny, J.P., 73
Cho, K., McFarlane, I.D., The anthozoan neuropeptide Antho-RWamide I modulates Ca^{2+} current in sea anemone myoepithelial cells, 53
Chung, J.M., see Lee, D.H., 140
Coalson, D.W., see Zacny, J.P., 73
Colling, S.B., Collinge, J., Jefferys, J.G.R., Hippocampal slices from prion protein null mice: disrupted Ca^{2+}-activated K^+ currents, 49
Collinge, J., see Colling, S.B., 49
Conneally, P.M., see Pritchard, M.L., 105
Courtois, F., see Giuliano, F., 1
Cullen, K.M., see Halliday, G.M., 29

D'Hooge, R., see Pan, J.C., 121
Damiani, D., see Carfagna, N., 189
De Deyn, P.P., see Pan, J.C., 121
Delogu, M.R., see Desole, M.S., 193
Desole, M.S., Sciola, L., Delogu, M.R., Sircana, S., Migheli, R., Manganese and 1-methyl-4-(2′-ethylphenyl)-1,2,3,6-tetrahydropyridine induce apoptosis in PC12 cells, 193
Dubbeldam, J.L., see Tellegen, A.J., 77

Eguchi, K., see Yokota, T., 204
Ekman, R., see Stiller, C.-O., 165

Fariello, R., see Carfagna, N., 189
Felgenhauer, K., see Blödorn, B., 117
Fergusson, J., see Layfield, R., 57
Feuerstein, T.J., Hüring, H., van Velthoven, V., Lücking, C.H., Landwehrmeyer, G.B., 5-HT_{1D}-like receptors inhibit the release of endogenously formed [^{3}H]GABA in human, but not in rabbit, neocortex, 210
Flamme, I., see Ikeda, E., 149
Franco, V.P., see Josselyn, S.A., 85

Gaskell, P.C., see Pritchard, M.L., 105
Gillin, J.C., see Seifritz, E., 41
Giuliano, F., Rampin, O., Brown, K., Courtois, F., Benoit, G., Jardin, A., Stimulation of the medial preoptic area of the hypothalamus in the rat elicits increases in intracavernous pressure, 1
Golanov, E.V., Reis, D.J., Cerebral cortical neurons with activity linked to central neurogenic spontaneous and evoked elevations in cerebral blood flow, 101
Griebel, R., see Kanthan, R., 207
Gundlach, A.L., see Burazin, T.C.D., 65

Haines, J.L., see Pritchard, M.L., 105
Hallbeck, M., Hermanson, O., Blomqvist, A., Preprovasopressin mRNA is not present in dorsal root ganglia of the rat, 125
Halliday, G.M., Cullen, K.M., Kril, J.J., Harding, A.J., Harasty, J., Glial fibrillary acidic protein (GFAP) immunohistochemistry in human cortex: a quantitative study using different antisera, 29
Harasty, J., see Halliday, G.M., 29
Harding, A.J., see Halliday, G.M., 29
Hayaishi, O., see Blödorn, B., 117
Hechler, V., see Cash, C.D., 25
Hermanson, O., see Hallbeck, M., 125
Herrero, J.F., Cervero, F., Supraspinal influences on the facilitation of rat nociceptive reflexes induced by carrageenan monoarthritis, 21
Hetta, J., see Imamura, K., 185
Higgins, M.J., Stone, T.W., Comparative sensitivity to adenosine of paired-pulse inhibition and single field potentials in the rat hippocampus, 69
Hirasawa, M., see Sakuragawa, N., 9
Hori, T., Suzuki, T., Baba, A., Abe, S., Yamamoto, T., Moroji, T., Shiraishi, H., Effects of phencyclidine metabolites on serotonin uptake in rat brain, 153
Horisaka, K., see Yamada, J., 113

Hui-Chan, C.W.Y., see Paquet, N., 215
Hüring, H., see Feuerstein, T.J., 210

Ikeda, E., Flamme, I., Risau, W., Developing brain cells produce factors capable of inducing the HT7 antigen, a blood–brain barrier-specific molecule, in chick endothelial cells, 149
Imamura, K., Onoe, H., Watanabe, Y., Andersson, J., Hetta, J., Schneider, H., Reineck, R., Lidström, K., Okura, M., Långström, B., Regional activation of human cerebral cortex upon an adaptation in mirror drawing, 185
Itoyama, Y., see Kondo, T., 157

Jardin, A., see Giuliano, F., 1
Jefferys, J.G.R., see Colling, S.B., 49
Jeppi, J.V., see Bulleit, R.F., 93
Josselyn, S.A., Franco, V.P., Vaccarino, F.J., PD-135158, a CCK_B receptor antagonist, microinjected into the nucleus accumbens and the expression of conditioned rewarded behaviour, 85

Kalra, J., see Kanthan, R., 207
Kamo, I., see Sakuragawa, N., 9
Kanthan, R., Shuaib, A., Griebel, R., Miyashita, H., Kalra, J., Glucose-induced decrease in glutamate levels in ischemic human brain by in-vivo microdialysis, 207
Kanzaki, R., see Okada, K., 197
Kase, H., see Kurokawa, M., 181
Kawachi, K., see Okada, K., 197
Kitai, S.T., see Plenz, D., 177
Kits, K.S., see Borst, J.G.G., 61
Klafta, J.M., see Zacny, J.P., 73
Klock, P.A., see Zacny, J.P., 73
Kondo, T., Shirasawa, T., Itoyama, Y., Mori, H., Embryonic genes expressed in Alzheimer's disease brains, 157
Kostopoulos, G.K., see Melkonian, D.S., 13
Kril, J.J., see Halliday, G.M., 29
Kurokawa, M., Shiozaki, S., Nonaka, H., Kase, H., Nakamura, J., Kuwana, Y., In vivo regulation of acetylcholine release via adenosine A_1 receptor in rat cerebral cortex, 181
Kuwana, Y., see Kurokawa, M., 181

Lai, L., see Pan, J.C., 121
Lambert, S., see Wintergerst, E.S., 173
Landon, M., see Layfield, R., 57
Landwehrmeyer, G.B., see Feuerstein, T.J., 210
Långström, B., see Imamura, K., 185
Layfield, R., Fergusson, J., Aitken, A., Lowe, J., Landon, M., Mayer, R.J., Neurofibrillary tangles of Alzheimer's disease brains contain 14-3-3 proteins, 57
Lee, D.H., Chung, J.M., Neuropathic pain in neonatal rats, 140
Leopold, E., see Mayer, B., 169
Lidström, K., see Imamura, K., 185
Lin, X., see Bulleit, R.F., 93
Lodder, J.C., see Borst, J.G.G., 61
Lowe, J., see Layfield, R., 57
Lücking, C.H., see Feuerstein, T.J., 210
Ludwig, M., see Papassotiropoulos, A., 33

Mäder, M., see Blödorn, B., 117
Maitre, M., see Cash, C.D., 25
Martin, D.L., see Battaglioli, G., 129
Matsumoto, M., see Yoshioka, M., 201
Mattammal, M.B., see Morley, J.E., 137
Mayer, B., Pfeiffer, S., Leopold, E., Müller, J., Weser, U., Schmidt, K., Structural and functional analogs of CuZn superoxide dismutase inhibit rat brain nitric oxide synthase by interference with the reductase (diaphorase) domain, 169
Mayer, R.J., see Layfield, R., 57
McFarlane, I.D., see Cho, K., 53
Melkonian, D.S., Kostopoulos, G.K., Novel approach to a quantitative treatment of the quantal transmitter release at the frog neuromuscular junction, 13
Mersel, M., see Cash, C.D., 25
Migheli, R., see Desole, M.S., 193
Miyashita, H., see Kanthan, R., 207
Mizuguchi, M., see Sakuragawa, N., 9
Mizuno, Y., see Mori, H., 109
Moore, P., see Seifritz, E., 41
Mori, H., Oda, M., Mizuno, Y., Cortical ballooned neurons in progressive supranuclear palsy, 109
Mori, H., see Kondo, T., 157
Morley, J.E., Mattammal, M.B., Nitric oxide synthase levels in obese Zucker rats, 137
Moroji, T., see Hori, T., 153
Müller, J., see Mayer, B., 169

Naib-Majani, W., see Papassotiropoulos, A., 33
Nakamura, J., see Kurokawa, M., 181
Nakamura, Y., see Otake, K., 97
Nonaka, H., see Kurokawa, M., 181
Norlén, M., Allard, P., Reduction in number of dopamine uptake sites but unchanged number of piperazine-acceptor/CYP450IID6 binding sites in the human caudate nucleus in aging, 161

Oda, M., see Mori, H., 109
Okada, K., Kanzaki, R., Kawachi, K., High-speed voltage-sensitive dye imaging of an in vivo insect brain, 197
Okura, M., see Imamura, K., 185
Onoe, H., see Imamura, K., 185
Otake, K., Nakamura, Y., Single neurons in Barrington's nucleus projecting to both the paraventricular thalamic nucleus and the spinal cord by way of axon collaterals: a double labeling study in the rat, 97

Pan, J.C., Pei, Y.Q., An, L., Lai, L., D'Hooge, R., De Deyn, P.P., Epileptiform activity and hippocampal damage produced by intrahippocampal injection of guanidinosuccinic acid in rat, 121
Papassotiropoulos, A., Ludwig, M., Naib-Majani, W., Rao, G.S., Induction of apoptosis and secondary necrosis in rat dorsal root ganglion cell cultures by oxidized low density lipoprotein, 33
Paquet, N., Tam, F., Hui-Chan, C.W.Y., Functional modulation of the human flexion and crossed extension reflexes by body position, 215
Pei, Y.Q., see Pan, J.C., 121
Pericak-Vance, M.A., see Pritchard, M.L., 105
Pfeiffer, S., see Mayer, B., 169
Phipps, B., see Piekut, D.T., 45
Piekut, D.T., Phipps, B., Corticotropin-releasing factor induction in rat piriform cortex following kainate-elicited seizures, 45
Plaghki, L., see Bragard, D., 81
Plenz, D., Kitai, S.T., Organotypic cortex-striatum-mesencephalon cultures: the nigrostriatal pathway, 177
Post, C., see Carfagna, N., 189
Pritchard, M.L., Saunders, A.M., Gaskell, P.C., Small, G.W.,

Conneally, P.M., Rosi, B., Yamaoka, L.H., Roses, A.D., Haines, J.L., Pericak-Vance, M.A., No association between very low density lipoprotein receptor (VLDL-R) and Alzheimer disease in American Caucasians, 105

Quercia, S., Chang, S.L., Antisense oligodeoxynucleotide attenuates in vivo expression of *c-fos* in the paraventricular hypothalamic nucleus of the rat brain, 89

Rampin, O., see Giuliano, F., 1
Rao, G.S., see Papassotiropoulos, A., 33
Rappelsberger, P., see Weiss, S., 17
Rathjen, F.G., see Wintergerst, E.S., 173
Reineck, R., see Imamura, K., 185
Reis, D.J., see Golanov, E.V., 101
Riederer, B.M., see Wintergerst, E.S., 173
Risau, W., see Ikeda, E., 149
Roses, A.D., see Pritchard, M.L., 105
Rosi, B., see Pritchard, M.L., 105
Roubos, E.W., see Borst, J.G.G., 61
Rupani, G., see Zacny, J.P., 73

Saito, H., see Yoshioka, M., 201
Sakuragawa, N., Thangavel, R., Mizuguchi, M., Hirasawa, M., Kamo, I., Expression of markers for both neuronal and glial cells in human amniotic epithelial cells, 9
Salmoiraghi, P., see Carfagna, N., 189
Satoh, T., see Yokota, T., 204
Saunders, A.M., see Pritchard, M.L., 105
Schmidt, K., see Mayer, B., 169
Schneider, H., see Imamura, K., 185
Sciola, L., see Desole, M.S., 193
Seifritz, E., Moore, P., Trachsel, L., Bhatti, T., Stahl, S.M., Gillin, J.C., The 5-HT_{1A} agonist ipsapirone enhances EEG slow wave activity in human sleep and produces a power spectrum similar to 5-HT_2 blockade, 41
Shiozaki, S., see Kurokawa, M., 181
Shiraishi, H., see Hori, T., 153
Shirasawa, T., see Kondo, T., 157
Shuaib, A., see Kanthan, R., 207
Sircana, S., see Desole, M.S., 193
Small, G.W., see Pritchard, M.L., 105
Stahl, S.M., see Seifritz, E., 41
Stiller, C.-O., Bergquist, J., Beck, O., Ekman, R., Brodin, E., Local administration of morphine decreases the extracellular level of GABA in the periaqueductal gray matter of freely moving rats, 165
Stone, T.W., see Higgins, M.J., 69
Strosznajder, J., Chalimoniuk, M., Strosznajder, R.P., Albanese, V., Alberghina, M., Arachidonate transport through the blood–retina and blood–brain barrier of the rat during aging, 145
Strosznajder, R.P., see Strosznajder, J., 145
Sugimoto, Y., see Yamada, J., 113
Sun, X., see Tian, X., 134
Suszkiw, J.B., see Tian, X., 134
Suzuki, T., see Hori, T., 153
Szente, M., see Boda, B., 37

Tam, F., see Paquet, N., 215
Tellegen, A.J., Dubbeldam, J.L., Do craniocervical and jaw motor nuclei receive input from the same population of reticular premotor neurons? a double labeling tracing study in the mallard (*Anas platyrhynchos*), 77
Teng, Y.D., Wrathall, J.R., Evaluation of cardiorespiratory parameters in rats after spinal cord trauma and treatment with NBQX, an antagonist of excitatory amino acid receptors, 5
Thangavel, R., see Sakuragawa, N., 9
Tian, X., Sun, X., Suszkiw, J.B., Developmental age-dependent upregulation of choline acetyltransferase and vesicular acetylcholine transporter mRNA expression in neonatal rat septum by nerve growth factor, 134
Togashi, H., see Yoshioka, M., 201
Trachsel, L., see Seifritz, E., 41

Urade, Y., see Blödorn, B., 117

Vaccarino, F.J., see Josselyn, S.A., 85
van Velthoven, V., see Feuerstein, T.J., 210
Vogt Weisenhorn, D.M., see Wintergerst, E.S., 173

Watanabe, Y., see Imamura, K., 185
Weiss, S., Rappelsberger, P., EEG coherence within the 13–18 Hz band as a correlate of a distinct lexical organisation of concrete and abstract nouns in humans, 17
Weser, U., see Mayer, B., 169
Wintergerst, E.S., Vogt Weisenhorn, D.M., Rathjen, F.G., Riederer, B.M., Lambert, S., Celio, M.R., Temporal and spatial appearance of the membrane cytoskeleton and perineuronal nets in the rat neocortex, 173
Wrathall, J.R., see Teng, Y.D., 5

Yamada, J., Sugimoto, Y., Yoshikawa, T., Horisaka, K., Effects of adrenodemedullation and adrenalectomy on the 5-HT_2 receptor agonists DOI- and mCPP-induced hypophagia in rats, 113
Yamamoto, T., see Hori, T., 153
Yamaoka, L.H., see Pritchard, M.L., 105
Yokota, T., Eguchi, K., Satoh, T., Correlated discharges of two neurons in rat gustatory cortex during gustatory stimulation, 204
Yoshikawa, T., see Yamada, J., 113
Yoshioka, M., Matsumoto, M., Togashi, H., Saito, H., Effect of conditioned fear stress on dopamine release in the rat prefrontal cortex, 201
Young, C.J., see Zacny, J.P., 73

Zacny, J.P., Cho, A.M., Coalson, D.W., Rupani, G., Young, C.J., Klafta, J.M., Klock, P.A., Apfelbaum, J.L., Differential acute tolerance development to effects of nitrous oxide in humans, 73